

Sacroiliac Joint Pain



PAIN MEDICINE ASSOCIATES
your life, your health, pain free

Long Beach Office
3620 Atlantic Ave Long Beach
Long Beach, CA 90807
(562) 595-0060 fax: (562) 595-0027

Fountain Valley Office
18111 Brookhurst St., Suite 3100
Fountain Valley, CA 90708
(714) 963-7240 fax: (714) 963-7224

www.PAINASSOCIATE.com

The sacroiliac joint (SI joint) is the joint bridging the lowest segment of the spinal column, the sacrum, to the bones of the pelvis. These joints support the upper body when we are standing. After time, the SI joint can develop arthritis and become painful, particularly when walking and climbing stairs. The location of pain is generally in the lower back, buttocks and thighs.

Imaging studies such as x-rays, MRI or CT may be unrevealing for SI joint pain and are typically obtained to rule out other lesions. Non-surgical treatment options are most appropriate to relieve SI joint pain. The first treatment options usually include chiropractic care, physical therapy and/or medication. If pain persists, injections may be done. Radiofrequency ablation, or rhizotomy, a procedure done to deaden nerves around the SI joint, may be performed to control pain.

1

DISCLAIMER: INFORMATION PROVIDED IN THIS SHEET IS GENERAL IN CONTENT AND SHOULD NOT BE SEEN AS A SUBSTITUTE FOR PROFESSIONAL MEDICAL ADVICE. CONCERNS OVER MEDICAL CONDITIONS SHOULD BE DISCUSSED WITH YOUR PRIMARY CARE PHYSICIAN OR PAIN CARE SPECIALIST.